

# Dynamical Neutrino Mass Matrix: Large (Small) Mixing Solution to Atmospheric (Solar) Neutrino Oscillations

Ernest Ma

*Department of Physics, University of California  
Riverside, California 92521, USA*

## Abstract

In the wake of the SuperKamiokande announcement of atmospheric neutrino oscillations which favor the large mixing of  $\nu_\mu$  with  $\nu_\tau$ , there is considerable theoretical activity towards a possible understanding of the solar neutrino deficit also in terms of the mixing of  $\nu_e$  with the other two neutrinos. A specific dynamical model is presented here for the first time which naturally chooses the large (small) mixing solution to atmospheric (solar) neutrino oscillations.

Separate experimental evidences of neutrino oscillations at three (or four) very different wavelengths have been accumulating for many years[1, 2, 3]. Recently the SuperKamiokande collaboration claimed[4] the strongest signal to date of atmospheric neutrino oscillations which favor the large mixing of  $\nu_\mu$  with  $\nu_\tau$ . Together with other known constraints, a simple emerging conclusion is that there exists a neutrino mass eigenstate  $\nu_\mu \cos \theta + \nu_\tau \sin \theta$  which is separated from the other states by a difference of mass squares ( $\Delta m^2$ ) of about  $10^{-3}$  to  $10^{-2}$  eV<sup>2</sup> and  $\sin^2 2\theta$  is greater than about 0.8. Within this context, if the solar neutrino deficit[1] is also to be explained, we have basically only three options, as depicted in Fig. 1.

In Option (a),  $\nu_e$  is the lowest lying state and mixes with  $c\nu_\tau - s\nu_\mu$ . If their  $\Delta m^2$  is from  $10^{-6}$  to  $10^{-5}$  eV<sup>2</sup>, then the matter-enhanced solution[5] to the solar neutrino deficit is possible. If their  $\Delta m^2$  is from  $10^{-11}$  to  $10^{-10}$  eV<sup>2</sup>, then the vacuum-oscillation solution[6] is possible. In Option (b),  $\nu_e$  is essentially degenerate with  $c\nu_\tau - s\nu_\mu$ , but it must still be below the latter to have the matter-enhanced solution. On the other hand, the vacuum-oscillation solution does not depend on the sign of  $\Delta m^2$  but requires near-maximal mixing between the two states to fit the data. In Option (c),  $\nu_e$  is placed below  $\nu_\mu$  with  $\Delta m^2$  of about 1 eV<sup>2</sup> to accommodate the LSND data[3]. In this case, a sterile neutrino  $\nu_s$  is needed just above  $\nu_e$  for the matter-enhanced solution to the solar neutrino deficit[7, 8]. This scenario of course also works without  $\nu_s$  if we were to discard the solar data. It should also be mentioned that  $\nu_e$  may still mix a little with  $c\nu_\mu + s\nu_\tau$ , as allowed by the SuperKamiokande experiment itself and within the constraint of the CHOOZ reactor data[9], as discussed already recently[10, 11].

To gain some insight into how the above-mentioned three options may be realized theoretically, let us look at each in turn. In Option (a), one mass is apparently much greater than the other two, whereas in Option (b), two nearly degenerate masses are much greater than the third. Of course, in either scenario, it is also possible to add a common mass to all three states without changing their relative standings. After all, neutrino oscillations only

measure  $\Delta m^2$  and not the individual masses. In Option (c), two nearly degenerate masses are much greater than two others. Note that whereas the gap between  $\nu_e$  and  $\nu_s$  is very small, the corresponding masses need not be degenerate in the sense that their ratio may in fact be very much less than unity.

If two neutrinos are nearly degenerate in mass, a natural explanation is that they are almost Dirac partners of each other. In that case, they mix maximally. In other words, there is a basis for which their  $2 \times 2$  mass matrix is approximately of the form

$$\mathcal{M}_\nu \simeq \begin{pmatrix} 0 & m \\ m & 0 \end{pmatrix}. \quad (1)$$

The mass eigenvalues of the above are  $\pm m$  and the mixing is  $45^\circ$ . The Majorana mass eigenstate which has the negative mass eigenvalue is redefined with a phase rotation, so that its mass becomes positive as required. In Option (c), this would automatically set  $c = s = 1/\sqrt{2}$  and we have a theoretical understanding[7] of why atmospheric neutrino oscillations favor  $\sin^2 2\theta = 1$ . In Option (b),  $\nu_e$  would mix maximally with  $c\nu_\tau - s\nu_\mu$  as proposed recently[10], and we have instead a theoretical understanding of why the vacuum-oscillation solution to the solar neutrino deficit favors  $\sin^2 2\theta = 1$ . In Option (a), it has also been assumed[12] that  $\nu_e$  mixes maximally with  $c\nu_\tau - s\nu_\mu$  although the two are not necessarily degenerate in the sense that their mass difference may not be small compared to the larger of the two masses. In both Options (a) and (b), the large mixing of  $\nu_\mu$  with  $\nu_\tau$ , which has the maximum value[13] of  $\sin^2 2\theta = 8/9$ , is not explained without further theoretical assumptions.

Most models of the neutrino mass matrix simply assume a certain pattern or texture, using whatever arguments the authors find to be attractive. A less arbitrary approach is to take the viewpoint that neutrino masses are in fact all zero at tree level, and ask how they can be generated dynamically in one loop. The first and simplest such model was due to Zee[14]. The dynamics of this model automatically tell us that  $\nu_\tau$  and a linear combination of  $\nu_\mu$  and  $\nu_e$

are almost Dirac partners. If the latter is dominantly  $\nu_\mu$ , then we have maximal atmospheric neutrino oscillations and the LSND results are also explained[15, 16]. To accommodate the solar data,  $\nu_s$  has to be added [Option (c)], but if its mass is also generated dynamically, then a proper extension[16] of the Zee model yields the interesting relationship

$$(\Delta m^2)_{\text{atm}} \simeq 2[(\Delta m^2)_{\text{solar}}(\Delta m^2)_{\text{LSND}}]^{1/2} \quad (2)$$

which is well satisfied by the data.

In this paper I consider instead a natural dynamical realization of Option (a), where the mixing of  $\nu_e$  with  $c\nu_\tau - s\nu_\mu$  is small, for the matter-enhanced solution[5] to the solar neutrino deficit. The object is to obtain a large mass for  $c\nu_\mu + s\nu_\tau$  and a small mass for  $c\nu_\tau - s\nu_\mu$ , with yet a smaller mass for  $\nu_e$ . To do this, I supplement the Zee mechanism with a global  $L_e$  symmetry, which is broken spontaneously as well as with explicit soft terms, by the interactions of the following new scalar particles.

$$(\eta^0, \eta^-), \chi^-, \zeta^0 : \quad L_e = 1. \quad (3)$$

Consider first the interactions given by

$$\mathcal{L}_1 = \sum_{i=\mu,\tau} f_i(\nu_i e - l_i \nu_e) \chi^+ + f'_i \bar{e}_R(\nu_i \eta^- - l_i \eta^0) + \mu_1(\phi^- \eta^0 + \bar{\phi}^0 \eta^-) \chi^+ + h.c., \quad (4)$$

where  $(\phi^+, \phi^0)$  is the standard Higgs doublet with  $L_e = 0$ . As shown in Fig. 2, the  $2 \times 2$  mass matrix spanning  $\nu_\mu$  and  $\nu_\tau$  is now of the form

$$m_{ij} = \frac{(f_i f'_j + f'_i f_j) \mu_1 v m_e}{16\pi^2 M^2}, \quad (5)$$

where  $v \equiv \langle \phi^0 \rangle = 174$  GeV and  $M$  is a large effective mass depending on  $m_\eta^2$  and  $m_\chi^2$ . Let  $f_1/f_2 \equiv \tan \alpha$  and  $f'_1/f'_2 \equiv \tan \beta$ , then the above can be rewritten as

$$m_{ij} = m_0 \begin{bmatrix} 2 \cos \alpha \cos \beta & \cos \alpha \sin \beta + \sin \alpha \cos \beta \\ \cos \alpha \sin \beta + \sin \alpha \cos \beta & 2 \sin \alpha \sin \beta \end{bmatrix}. \quad (6)$$

The trace of  $m_{ij}$  is  $2m_0 \cos(\alpha - \beta)$  and its determinant is  $-m_0^2 \sin^2(\alpha - \beta)$ .

Since  $\tan \alpha$  and  $\tan \beta$  are naturally of order unity, a very plausible scenario is to have  $\cos(\alpha - \beta) \sim 1$  and  $\sin^2(\alpha - \beta) \ll 1$ . In that case, the mass eigenvalues are approximately  $2m_0$  and  $-(m_0/2) \sin^2(\alpha - \beta)$ , corresponding to the eigenstates  $c\nu_\mu + s\nu_\tau$  and  $c\nu_\tau - s\nu_\mu$  respectively, where

$$\frac{s}{c} \simeq \frac{\sin(\alpha + \beta)}{2 \cos \alpha \cos \beta}, \quad (7)$$

which is equal to one for  $\alpha = \beta = 45^\circ$ . Numerically, let  $2m_0 \simeq 0.07$  eV, then the other mass would be  $2.3 \times 10^{-3}$  eV for  $\sin^2(\alpha - \beta) \simeq 0.13$ . These are very reasonable numbers and they are phenomenologically very relevant because they give  $\Delta m^2 \simeq 5 \times 10^{-3}$  eV<sup>2</sup> for atmospheric neutrino oscillations and  $\Delta m^2 \simeq 5 \times 10^{-6}$  eV<sup>2</sup> for solar neutrino oscillations. To obtain  $2m_0 \simeq 0.07$  eV from Eq. (5), we need  $ff'\mu_1/M^2 \simeq 6.2 \times 10^{-8}$  GeV<sup>-1</sup>, which may be achieved with  $f = f' = 0.05$ ,  $\mu_1 = 100$  GeV, and  $M = 2$  TeV.

So far, the one-loop diagram of Fig. 2 has yielded the following mass matrix in the basis  $(\nu_e, c\nu_\tau - s\nu_\mu, c\nu_\mu + s\nu_\tau)$ :

$$\mathcal{M}_\nu = \begin{bmatrix} 0 & 0 & 0 \\ 0 & m_2 & 0 \\ 0 & 0 & m_3 \end{bmatrix}, \quad (8)$$

where  $m_2 \simeq 2.3 \times 10^{-3}$  eV and  $m_3 \simeq 0.07$  eV. To have matter-enhanced solar neutrino oscillations,  $\nu_e$  must be made to mix with  $c\nu_\tau - s\nu_\mu$  and its mass must be smaller than  $m_2$ . To accomplish this, the neutral singlet  $\zeta^0$  of Eq. (3) is now used. It contributes to the interactions given by

$$\mathcal{L}_2 = \mu_2(\phi^+\eta^- - \phi^0\eta^0)\bar{\zeta}^0 + \mu_3(\phi^+\phi^- + \phi^0\bar{\phi}^0)\zeta^0 + \mu_4(\eta^+\eta^- + \bar{\eta}^0\eta^0)\zeta^0 + h.c. \quad (9)$$

Whereas the  $\mu_2$  term conserves  $L_e$ , the others do not. Hence  $L_e$  is softly broken explicitly. Furthermore, as  $\phi^0$  acquires a nonzero vacuum expectation value ( $vev$ ), so must  $\zeta^0$  and  $\eta^0$ . Hence  $L_e$  is also broken spontaneously, but there is no massless Goldstone boson because

$L_e$  is already broken explicitly. A nonzero  $\langle \eta^0 \rangle$  means that the charged-lepton mass matrix picks up nondiagonal terms from Eq. (4). The  $3 \times 3$  matrix linking  $(\bar{e}_L, \bar{\mu}_L, \bar{\tau}_L)$  to  $(e_R, \mu_R, \tau_R)$  becomes

$$\mathcal{M}_l = \begin{bmatrix} m_e & 0 & 0 \\ f'_\mu \langle \eta^0 \rangle & m_\mu & 0 \\ f'_\tau \langle \eta^0 \rangle & 0 & m_\tau \end{bmatrix}, \quad (10)$$

and the re-diagonalization of this matrix induces a mixing of  $\nu_e$  with  $\nu_\mu$  and  $\nu_\tau$  of magnitude  $f'_\mu \langle \eta^0 \rangle m_e / m_\mu^2$  and  $f'_\tau \langle \eta^0 \rangle m_e / m_\tau^2$  respectively. The former should be much greater than the latter and it may not be negligible if  $\langle \eta^0 \rangle$  is not small. However, if  $m_\eta^2$  is positive and large, then

$$\langle \eta^0 \rangle \simeq \frac{\mu_2 v \langle \zeta^0 \rangle}{m_\eta^2}. \quad (11)$$

In other words, a heavy  $\eta$  may have a naturally small  $vev$ , as for example shown recently [17].

In the presence of a nonzero  $\langle \eta^0 \rangle$ , there is also a dynamical mass linking  $\nu_e$  with  $\nu_\mu$  and  $\nu_\tau$  as shown in Fig. (3), as well as one with itself as shown in Fig. 4. The  $3 \times 3$  matrix of Eq. (8) is now changed:

$$\mathcal{M}_\nu = \begin{bmatrix} m'' & cm' & sm' \\ cm' & m_2 & 0 \\ sm' & 0 & m_3 \end{bmatrix}, \quad (12)$$

where

$$m' = \frac{f_\tau m_\tau^2 \mu_1 \langle \eta^0 \rangle}{16\pi^2 v m_\chi^2}, \quad m'' = \frac{(f_\mu f'_\mu + f_\tau f'_\tau) m_e \mu_1 \langle \eta^0 \rangle^2}{16\pi^2 v m_\chi^2}. \quad (13)$$

Let  $\langle \eta^0 \rangle = 0.9$  MeV, and using the same values of  $f = f' = 0.05$ ,  $\mu_1 = 100$  GeV and  $m_\chi = 2$  TeV as before,  $m'$  is then  $1.3 \times 10^{-4}$  eV and  $m''$  is  $1.9 \times 10^{-12}$  eV. Assuming  $c \simeq 1/\sqrt{2}$ , the mixing of  $\nu_e$  with  $c\nu_\tau - s\nu_\mu$  in Eq. (12) is

$$\frac{cm'}{m_2} = 0.04, \quad (14)$$

which yields the value of  $\sin^2 2\theta = 6 \times 10^{-3}$  for solar neutrino oscillations.

Going back to Eq. (10), it is easy to check that the value of  $\langle \eta^0 \rangle = 0.9$  MeV yields a mixing of  $2 \times 10^{-6}$  for  $\nu_e$  with  $\nu_\mu$  which is clearly negligible. From Eq. (11), using again

$m_\eta = 2 \text{ TeV}$ , it is seen that  $\mu_2 \langle \zeta^0 \rangle = 21 \text{ GeV}^2$ , which is very reasonable. The mass of the state dominated by  $\nu_e$  is

$$m_1 = m'^2/2m_2 = 3.7 \times 10^{-6} \text{ eV}, \quad (15)$$

the contribution  $m''$  from Fig. 4 being negligible.

In summary, it has been shown in this paper for the first time how a specific dynamical model naturally chooses the large (small) mixing solution to atmospheric (solar) neutrino oscillations. Unlike most other attempts to understand the present data, this is a complete theoretical model and not just an arbitrary speculation on the form of the neutrino mass matrix. The three approximate mass eigenstates are  $\nu_\mu \cos \theta + \nu_\tau \sin \theta$ ,  $\nu_\tau \cos \theta - \nu_\mu \sin \theta$  and  $\nu_e$ , with eigenvalues  $0.07 \text{ eV}$ ,  $2.3 \times 10^{-3} \text{ eV}$  and  $3.7 \times 10^{-6} \text{ eV}$  respectively. The  $\nu_\mu - \nu_\tau$  mixing angle  $\theta$  is naturally large and could be close to  $45^\circ$ , thus accounting for the atmospheric data[2, 4]. The mixing of  $\nu_e$  with  $\nu_\tau \cos \theta - \nu_\mu \sin \theta$  is about 0.04, resulting in a value of  $\sin^2 2\theta = 6 \times 10^{-6}$ , which is a suitable matter-enhanced solution to the solar data[1].

The  $3 \times 3$  neutrino mass matrix, as given by Eq. (12), is obtained dynamically in one loop, as depicted in Figs. 2 to 4. This is achieved with a minimal set of new heavy scalar particles, as listed in Eq. (3), which have  $L_e = 1$ , *i.e.* electron number just as  $\nu_e$  and  $e$ . In the limit of exact  $L_e$  conservation, the mass matrix of Eq. (8) is obtained. This is sufficient to explain the atmospheric data. As  $L_e$  is broken spontaneously (and with explicit soft terms so that a massless Goldstone boson will not appear),  $\nu_e$  mixes with  $\nu_\tau$  and matter-enhanced solar neutrino oscillations become possible. This requires a small  $\langle \eta^0 \rangle$ , which is actually very natural[17] for a positive and large  $m_\eta^2$ . The above scenario [Option(a) of Fig. 1] is applicable as long as the LSND data[3] are not considered. If all positive neutrino-oscillation data are to be accommodated, an extension of the Zee model[14] to include a sterile neutrino [Option(c)] could be the answer[16]. In either case, a dynamical explanation of the neutrino mass matrix works very well in the present experimental context.

## ACKNOWLEDGEMENT


This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

## References


- [1] R. Davis, Prog. Part. Nucl. Phys. **32**, 13 (1994); Y. Fukuda *et al.*, Phys. Rev. Lett. **77**, 1683 (1996); hep-ex/9805021; P. Anselmann *et al.*, Phys. Lett. **B357**, 237 (1995); **B361**, 235 (1996); J. N. Abdurashitov *et al.*, Phys. Lett. **B328**, 234 (1994).
- [2] D. Casper *et al.*, Phys. Rev. Lett. **66**, 2561 (1991); R. Becker-Szendy *et al.*, Phys. Rev. **D46**, 3720 (1992); K. S. Hirata *et al.*, Phys. Lett. **B280**, 146 (1992); Y. Fukuda *et al.*, Phys. Lett. **B335**, 237 (1994); S. Hatakeyama *et al.*, hep-ex/9806038.
- [3] C. Athanassopoulos *et al.*, Phys. Rev. Lett. **75**, 2650 (1995); **77**, 3082 (1996); nucl-ex/9709006 (Phys. Rev. Lett., in press).
- [4] Y. Fukuda *et al.*, hep-ex/9803006,9805006,9807003; T. Kajita, in Proceedings of the XVIIIth International Conference on Neutrino Physics and Astrophysics, Takayama, Japan ( June 1998).
- [5] L. Wolfenstein, Phys. Rev. **D17**, 2369 (1978); **20**, 2364 (1979); S. Mikheyev and A. Yu. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985).
- [6] V. Barger, R. J. N. Phillips, and K. Whisnant, Phys. Rev. **D24**, 538 (1981); S. L. Glashow and L. M. Krauss, Phys. Lett. **B190**, 199 (1987).
- [7] E. Ma and P. Roy, Phys. Rev. **D52**, R4780 (1995).




- [8] N. Okada and O. Yasuda, Int. J. Phys. **A12**, 3669 (1997); S. M. Bilenky, C. Giunti, and W. Grimus, Eur. Phys. **C1**, 247 (1998).
- [9] M. Apollonio *et al.*, Phys. Lett. **B420**, 397 (1998).
- [10] R. Barbieri, L. J. Hall, D. Smith, A. Strumia, and N. Weiner, hep-ph/9807235.
- [11] V. Barger, T. Weiler, and K. Whisnant, hep-ph/9807319.
- [12] V. Barger, S. Pakvasa, T. J. Weiler, and K. Whisnant, hep-ph/9806387; A. J. Baltz, A. S. Goldhaber, and M. Goldhaber, hep-ph/9806540; M. Tanimoto, hep-ph/9807283; Y. Nomura and T. Yanagida, hep-ph/9807325.
- [13] H. Fritzsch and Z. Xing, Phys. Lett. **B372**, 265 (1996); M. Fukugita, M. Tanimoto, and T. Yanagida, Phys. Rev. **D57**, 4429 (1998).
- [14] A. Zee, Phys. Lett. **93B**, 389 (1980).
- [15] A. Yu. Smirnov and M. Tanimoto, Phys. Rev. **D55**, 1665 (1997).
- [16] N. Gaur, A. Ghosal, E. Ma, and P. Roy, hep-ph/9806272.
- [17] E. Ma and U. Sarkar, Phys. Rev. Lett. **80**, 5716 (1998).




**Fig. 1.** Schematic drawings of possible neutrino mass hierarchies.



**Fig. 2.** Dynamical mass generation for  $\nu_\mu$  and  $\nu_\tau$ .



**Fig. 3.** Dynamical mass linking  $\nu_e$  with  $\nu_\tau$ .



**Fig. 4.** Dynamical mass for  $\nu_e$ .